Brain Research, 412 (1987) 409–411
Elsevier

Author Index

Abe, K., see Tamamaki, N., 156
Aravich, P.F., see Silverman, W.F., 375
Armstrong, R., Toews, A.D. and Morell, P., Axonal transport through nodes of Ranvier, 196

Badoer, E., see Head, G.A., 18
Barbeau, H. and Rossignol, S., Recovery of locomotion after chronic spinalization in the adult cat, 84
Barbin, G., see Simmonneau, M., 224
Benassi, C., see Biral, G.P., 43
Benetos, A., Gavras, I. and Gavras, H., Stimulation of vasopressin by calcium microinjections in the area of the paraventricular nucleus of the hypothalamus, 182
Beninato, M. and Spencer, R.F., A cholinergic projection to the rat substantia nigra from the pedunculopontine tegmental nucleus, 169
Bhat, S. and Silberberg, D.H., C6 glioma cells express modified neural-cell adhesion molecule-like glycoproteins, 144
Biral, G.P., Porro, C.A., Cavazzuti, M., Benassi, C. and Corazza, R., Vertical and horizontal visual whole-field motion differently affect the metabolic activity of the rat medial terminal nucleus, 43
Bittiger, H., see Olpe, H.R., 269
Black, I.B., see Martínez, H.J., 295
Bonhaus, D.W., see Shin, C., 311
Boulton, A.A., see Sardar, A.A., 370
Bourgoin, S., see Le Bars, D., 190
Brecha, N.C., see Mantyh, C.R., 329
Bulloch, A.G.M., Somatostatin enhances neurite outgrowth and electrical coupling of regenerating neurons in *Helisoma*, 6
Busto, R., see Yoshida, S., 114

Cavazzuti, M., see Biral, G.P., 43
Cesselin, F., see Le Bars, D., 190
Chang, K.-J., see Crain, B.J., 343
Changaris, D.G., see Schurr, A., 179
Church, W.H., Justice Jr., J.B. and Neill, D.B., Detecting behaviorally relevant changes in extracellular dopamine with microdialysis, 397
Clot, A.M., see Le Bars, D., 190
Collier, T.J., see Silverman, W.F., 375
Corazza, R., see Biral, G.P., 43
Cottrell, G.A. and Robertson, H.A., Prevention of cysteamine-induced myoclonus blocks the long-term inhibition of kindled seizures, 161
Crain, B.J., Chang, K.-J. and McNamara, J.O., An in vitro autoradiographic analysis of mu and delta opioid binding in the hippocampal formation of kindled rats, 343
Cynader, M.S., see Prusky, G.T., 131

Davidsson, P., Karlsson, B. and Svennerholm, L., Glycoprotein pattern in human brain tumors studied using lectin binding after sodium dodecyl sulfate-gel electrophoresis and protein blotting, 254
De Jonge, M. and Racine, R.J., The development and decay of kindling-induced increases in paired-pulse depression in the dentate gyrus, 318
Denton, D.A., see Osborne, P.G., 36
Distasi, C., see Simmonneau, M., 224
Drachman, D.B., see Pestronk, A., 302
Dreyfus, C.F., see Martínez, H.J., 295

Forman, D.S., Lynch, K.J. and Smith, R.S., Organelle dynamics in lobster axons: anterograde, retrograde and stationary mitochondria, 96
Fuchs, I., see Sanghera, M.K., 200
Fujishima, M., see Yoshida, F., 1
Fukui, M., see Kondo, A., 73

Gavras, H., see Benetos, A., 182
Gavras, I., see Benetos, A., 182
Gentile, C.G., see Jarrell, T.W., 285
Gerard, P., see Huguet, F., 125
Ginsberg, M.D., see Yoshida, S., 114
Gola, M., see Pin, T., 165
Gonzalez-Lima, F., Midbrain reticular stimulation produces patterns of metabolic activation and suppression in the cerebellum and vestibular nuclei: a 2-deoxyglucose study, 275
González, J.L. and Santos-Benito, F.F., Synthesis of acetylcholine by endothelial cells isolated from rat brain cortex capillaries, 148
Graziadei, P.P.C., see Magrassi, L., 386
Groenewegen, H.J., see Voorn, P., 391

Hamon, M., see Le Bars, D., 190
Hamprecht, B., see Höpp, H.-P., 175
Hatakenaka, S., see Pasteels, B., 107
Hayes, K.C., see Sullivan, S.J., 139
Head, G.A., Badoer, E. and Korner, P.I., Cardiovascular role of A_1 catecholaminergic neurons in the rabbit. Effect of chronic lesions on responses to methyldopa, clonidine and 6-OHDA induced transmitter release, 18
Heid, J., see Olpe, H.R., 269
Henderson, Z., Source of cholinergic input to ferret visual cortex, 261
Henderson, Z., A small proportion of cholinergic neurones in the nucleus basalis magnocellularis of ferret appear to stain positively for tyrosine hydroxylase, 363
Hong, J.S., see Sivam, S.P., 29
Huchet, A.-M., see Huguet, F., 125
Hudson, P.M., see Sivam, S.P., 29
Huguet, F., Huchet, A.-M., Gerard, P. and Narcisse, G., Characterization of dihydropyridine binding sites in the rat brain: hypertension and age-dependent modulation of [^{3}H](+)-PN 200-110 binding, 125
Hunt, S.P., see Waters, C.M., 244
Höpp, H.-P., Reuter, G., Reiser, G. and Hamprecht, B., Angiotensin evokes in polyploid rat glioma cells hyperpolarization–depolarization responses and cross-desensitization with bradykinin, 175

Ikeda, M., see Yoshida, S., 114
Inoue, T., see Kondo, A., 73

Ishituka, T., see Yoshida, F., 1

Jarrell, T.W., Gentile, C.G., Romanski, L.M., McCabe, P.M. and Schneiderman, N., Involvement of cortical and thalamic auditory regions in retention of differential bradycardiac conditioning to acoustic conditioned stimuli in rabbits, 285
Jeneskog, T., Termination in posterior and anterior cerebellum of a climbing fibre pathway activated from the nucleus of Darkschewitsch in the cat, 185
Jenner, P., see Waters, C.M., 244
Jonakait, G.M., see Martínez, H.J., 295
Joyce, J.N., see Rhodes, K.J., 400
Juorio, A.V., see Sardar, A., 370
Justice Jr., J.B., see Church, W.H., 397

Karlsson, B., see Davidsson, P., 254
Khayutin, V.M., see Lukoshkova, E.V., 357
Kocsis, B., see Lukoshkova, E.V., 357
Kondo, A., Inoue, T., Nagara, H., Tateishi, J. and Fukui, M., Neurotoxicity of adriamycin passed through the transiently disrupted blood–brain barrier by mannitol in the rat brain, 73
Koob, G.F., see Swerdlow, N.R., 233
Korner, P.I., see Head, G.A., 18
Kruger, L., see Mantyh, C.R., 329

Lauritzen, M., see Okada, Y.C., 151
Le Bars, D., Bourgoin, S., Villanueva, L., Clot, A.M., Hamon, M. and Cesselin, F., Involvement of the dorsolateral funiculi in the spinal release of Met-enkephalin-like material triggered by heterosegmental noxious mechanical stimuli, 190
Lukoshkova, E.V., Kocsis, B., Shuvalova, T.B. and Khayutin, V.M., Characterization of the very late cardiosympathetic A-reflex in high-mesencephalic cats, 357
Lynch, K.J., see Forman, D.S., 96

Magrassi, L. and Graziadei, P.P.C., Single olfactory organ associated with prosencephalic malformation and cyclopia in a *Xenopus laevis* tadpole, 386
Mantyh, C.R., Kruger, L., Brecha, N.C. and Mantyh, P.W., Localization of specific binding sites for atrial natriuretic factor in the central nervous system of rat, guinea pig, cat and human, 329
Mantyh, P.W., see Mantyh, C.R., 329
Marsden, C.D., see Waters, C.M., 244
Marshall, J.F., see Rhodes, K.J., 400
Martinez, E., see Yoshida, S., 114
Martínez, H.J., Dreyfus, C.F., Jonakait, G.M. and Black, I.B., Nerve growth factor selectively increases cholinergic markers but not neuropeptides in rat basal forebrain in culture, 295
McCabe, P.M., see Jarrell, T.W., 285
McLaughlin, P.J., see Zagon, I.S., 68
McNamara, J.O., see Crain, B.J., 343
McNamara, J.O., see Shin, C., 311
Miceli, M.O., Post, C.A. and Van der Kooy, D., Catecholamine and serotonin colocalization in projection neurons of the area postrema, 381
Miki, N., see Pasteels, B., 107
Morell, P., see Armstrong, R., 196
Mozaffari, B., see Walz, W., 405

Nagara, H., see Kondo, A., 73
Narcisse, G., see Huguet, F., 125
Neal, J.W., see Pearson, R.C.A., 312
Neill, D.B., see Church, W.H., 397
Nicholson, C., see Okada, Y.C., 151
Nojyo, Y., see Tamamaki, N., 156

Ogata, J., see Yoshida, F., 1
Okada, Y.C., Lauritzen, M. and Nicholson, C., Magnetic field associated with neural activities in an isolated cerebellum, 151
Olpe, H.R., Heid, J., Bittiger, H. and Steinmann, M.W., Substance P depresses neuronal activity in the rat olfactory bulb in vitro and in vivo: possible mediation via γ-aminobutyric acid release, 269
Olschowka, J.A., see Silverman, W.F., 375
Osborne, P.G., Denton, D.A. and Weisinger, R.S., Inhibition of dehydration induced drinking in rats by reduction of CSF Na concentration, 36

Pasteels, B., Miki, N., Hatakenaka, S. and Pochet, R., Immunohistochemical cross-reactivity and electrophoretic comigration between calbindin D-27 kDa and visinin, 107
Pearson, R.C.A., Neal, J.W. and Powell, T.P.S., Increase in immunohistochemical staining of GABAergic axons in the superior colliculus and thalamus of the rat following damage of the ipsilateral striatum and frontal cortex, 312
Pestronk, A. and Drachman, D.B., Mechanism of action of lithium on acetylcholine receptor metabolism in skeletal muscle, 302
Pin, T. and Gola, M., Decreased resting membrane conductance as the proximate cause of burst modulation by an identified interneuron in *Helix*, 165
Pochet, R., see Pasteels, B., 107
Porro, C.A., see Biral, G.P., 43
Porter, L.H., An experimental investigation of the parietal lobes and temperature discrimination in monkeys, 54
Post, C.A., see Miceli, M.O., 381
Powell, T.P.S., see Pearson, R.C.A., 312
Prusky, G.T., Shaw, C. and Cynader, M.S., Nicotine receptors are located on lateral geniculate nucleus terminals in cat visual cortex, 131

Racine, R.J., see De Jonge, M., 318
Reiser, G., see Höpp, H.-P., 175
Reuter, G., see Höpp, H.-P., 175
Rhodes, K.J., Joyce, J.N., Sapp, D.W. and Marshall, J.F., [^{3}H]Hemicholinium-3 binding in rabbit striatum: correspondence with patchy acetylcholinesterase staining and a method for quantifying striatal compartments, 400
Rigor, B.M., see Schurr, A., 179
Robertson, H.A., see Cottrell, G.A., 161
Roest, G., see Voorn, P., 391
Romanski, L.M., see Jarrell, T.W., 285
Rossignol, S., see Barbeau, H., 84

Sadoshima, S., see Yoshida, F., 1
Sanghera, M.K., Fuchs, I., Weidmer-Mikhail, E. and Speciale, S.G., Met-Enkephalin levels in midbrain dopamine regions of inbred mouse strains which differ in the number of dopamine neurons, 200
Santiso, M., see Yoshida, S., 114
Santos-Benito, F.F., see González, J.L., 148
Sapp, D.W., see Rhodes, K.J., 400
Sardar, A., Juorio, A.V. and Boulton, A.A., The concentra-

tion of *p*- and *m*-tyramine in the rat mesolimbic system: its regional distribution and effect of monoamine oxidase inhibition, 370
Schneiderman, N., see Jarrell, T.W., 285
Schurr, A., Changaris, D.G. and Rigor, B.M., Glutamine protects neuronal function against cerebral hypoxia: a study using the in vitro hippocampal slice preparation, 179
Shaw, C., see Prusky, G.T., 131
Shin, C., Silver, J.M., Bonhaus, D.W. and McNamara, J.O., The role of substantia nigra in the development of kindling: pharmacologic and lesion studies, 311
Shuvalova, T.B., see Lukoshkova, E.V., 357
Silberberg, D.H., see Bhat, S., 144
Silver, J.M., see Shin, C., 311
Silverman, W.F., Aravich, P.F., Collier, T.J., Olschowka, J.A. and Sladek Jr., J.R., Reinnervation of transplanted hypothalamic neurons by host aminergic fibers in rats, 375
Simmonneau, M., Distasi, C., Tauc, L. and Barbin, G., Potassium channels in mouse neonate dorsal root ganglion cells: a patch-clamp study, 224
Sivam, S.P., Hudson, P.M., Tilson, H.A. and Hong, J.S., GABA and dopamine interaction in the basal ganglia: dopaminergic supersensitivity following chronic elevation of brain γ-aminobutyric acid levels, 29
Sladek Jr., J.R., see Silverman, W.F., 375
Smith, R.S., see Forman, D.S., 96
Speciale, S.G., see Sanghera, M.K., 200
Spencer, R.F., see Beninato, M., 169
Steinmann, M.W., see Olpe, H.R., 269
Sullivan, S.J. and Hayes, K.C., Changes in short and long latency stretch reflexes prior to movement initiation, 139
Svennerholm, L., see Davidsson, P., 254
Swerdlow, N.R. and Koob, G.F., Lesions of the dorsomedial nucleus of the thalamus, medial prefrontal cortex and pedunculopontine nucleus: effects on locomotor activity mediated by nucleus accumbens-ventral pallidal circuitry, 233

Tamamaki, N., Abe, K. and Nojyo, Y., Columnar organization in the subiculum formed by axon branches originating from single CA_1 pyramidal neurons in the rat hippocampus, 156
Tateishi, J., see Kondo, A., 73
Tauc, L., see Simmonneau, M., 224
Tilson, H.A., see Sivam, S.P., 29
Toews, A.D., see Armstrong, R. 196

Van der Kooy, D., see Miceli, M.O., 381
Villanueva, L., see Le Bars, D., 190
Voorn, P., Roest, G. and Groenewegen, H.J., Increase of enkephalin and decrease of substance P immunoreactivity in the dorsal and ventral striatum of the rat after midbrain 6-hydroxydopamine lesions, 391

Walz, W. and Mozaffari, B., Culture environment and channel-mediated potassium fluxes in astrocytes, 405
Waters, C.M., Hunt, S.P., Jenner, P. and Marsden, C.D., Localization of neurotensin receptors in the forebrain of the common marmoset and the effects of treatment with the neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine, 244
Weidmer-Mikhail, E., see Sanghera, M.K., 200
Weisinger, R.S., see Osborne, P.G., 36
Wild, J.M., The avian somatosensory system: connections of regions of body representation in the forebrain of the pigeon, 205

Yoshida, F., Fujishima, M., Sadoshima, S., Ishituka, T. and Ogata, J., Carbon dioxide reactivity of cerebral cortical and pial arteries in spontaneously hypertensive and normotensive rats — a morphometric study, 1
Yoshida, S., Ikeda, M., Busto, R., Santiso, M., Martinez, E. and Ginsberg, M.D., Cerebral phosphoinositide, triacylglycerol and energy metabolism during sustained seizures induced by bicuculline, 114

Zagon, I.S. and McLaughlin, P.J., Endogenous opioid systems regulate cell proliferation in the developing rat brain, 68